

SYSTEMATIC EFFECTS IN THE INTERPRETATIONS OF CLUSTER X-RAY TEMPERATURE FUNCTIONS

S. M. Molnar ^{1,2} and K. Jahoda ¹

*1) Laboratory for High Energy Astrophysics, Code 662 Goddard Space Flight Center
Greenbelt, MD 20771*

2) NAS/NRC Research Associate

ABSTRACT The formation and evolution of clusters of galaxies are sensitive to the underlying cosmological model. Constraints on cosmological parameters of cold dark matter models have been derived from mass, temperature and luminosity functions of clusters. We study the importance of including cluster formation history and a correction for collapsed fraction of objects in determining the cluster X-ray temperature function. We find that both effects are important. We compare temperature functions obtained by using a power law approximation for the mass variance normalized to X-ray clusters to those obtained by using a COBE normalized full CDM treatment. We conclude that the temperature function could be a powerful test on the average density of the Universe if we could find the correct way of interpreting the data.

KEYWORDS: galaxies: clusters: general — large-scale structure of universe.

1. INTRODUCTION

Clusters of galaxies are the largest known virialized gravitational systems in the Universe. According to our working hypothesis of structure formation, the cold dark matter (CDM) model, clusters form via gravitational collapse of high density peaks of primordial density fluctuations. The distribution and evolution of these peaks strongly depend on the CDM model parameters. We use the following parameterization for our CDM models: Ω_0 , Ω_b , Ω_Λ , h (matter density, baryon density, cosmological constant, all in units of the critical density, all at $z = 0$, and h is the Hubble constant in units of $100 \text{ Kms}^{-1} \text{ Mpc}^{-1}$). Semi-analytical methods, like the Press-Schechter approximation, use some version of a spherical collapse model to predict the distribution of collapsed objects and their masses as a function of redshift. Assuming a physical model for the clusters and their intracluster medium, one can translate cluster virial masses to X-ray luminosity, or temperature of the intracluster gas, which can be compared to observations. We use the temperature function of clusters in our study. The most popular method is to assume a power law approximation for the power spectra of CDM models, use the Press-Schechter mass function (PSMF) to predict the abundance, and then derive the virial temperature to get the intracluster gas temperature, or integrate over formation epochs to

determine the mass-temperature conversion. This method has the power spectrum normalization and the power law exponent of the mass variance on cluster scales as free parameters and determines matter density, and has no dependence on the Hubble constant. We use COBE normalized spectra of CDM models to derive the mass variance and the Press-Schechter mass function to derive the distribution of cluster masses. This method returns the matter density as a function of the Hubble constant, thus one can not derive either parameter separately. We use four different methods to obtain theoretical temperature functions. We integrate over formation epochs in methods A and B but not in methods C and D. We take into account the collapsed fraction of objects in methods A and C and not in B and D. Method A thus take both effects and method D none of them into account. For each method (A, B, C, and D) we use a grid of CDM model parameters, $0.2 < \Omega_m < 0.9$ and $0.2 < h < 0.9$, and calculate the temperature functions assuming open and flat CDM cosmologies. We compare these theoretical temperature functions to data and determine the best fit models by minimizing the corresponding χ^2 .

2. OUTLINE OF THE METHODS

The power law approximation for the mass variance assumes a power exponent α , and derives the mass variance as $\sigma(M) = \sigma_8(M/M_{15})^{-\alpha}$, where $M_{15} = M/(10^{15} M_\odot)$, and M is the mass we will identify with the virial mass in the PSMF. It is commonly assumed that the power spectrum then can be approximated by a power law with an exponent $\alpha = (n_{PS} + 3)/6$, but that is true only if the power spectrum could be approximated by one power law in all scales (note the integral in equation 1). That is not true for CDM models, therefore we quote the mass variance power exponents on cluster scale for power law approximations. Instead of the approximation we use COBE normalized power spectra of CDM models (Hu and Sugiyama 1996) and obtain the mass variance from a numerical integral

$$\sigma_{R(M)}^2 = (1/2\pi^2) \int P(k) W^2(kR) k^2 dk, \quad (1)$$

where $P(k)$, $W(kR)$, $R(M)$, and k are the power spectrum, filter function in Fourier space, the radius of filtering, and the co-moving wavenumber. We use the standard PSMF corrected for collapsed fraction f_c , which gives the fraction of matter in collapsed objects: $n(z, M) dM = (f_c/f_c^{PS}) n_{PS}(z, M) dM$, where f_c^{PS} is the Press-Schechter collapsed fraction (Martel and Shapiro 1999). In order to obtain the temperature function, we assume that clusters form over a period of time and integrate over formation $n_T(z, M) = \int_z^\infty F(M, z_f, z) \frac{dM}{dT}(T, z_f, z) dz_f$, where $F(M, z_f, z)$ depends on $\sigma(M)$, $\delta_c(z)$, and $n(z_f, M)$ (Kitayama and Suto 1996). We used the spherical collapse model virial temperature (Eke et al. 1998):

$$M(T) = M_{15} \left[\frac{\beta k T [keV]}{1.38 keV (z + 1)} \right]^{\frac{3}{2}} \left[\frac{\Omega(z)}{\Omega(0) \Delta_c(z)} \right]^{\frac{1}{2}}. \quad (2)$$

We use $\beta = 1$ as suggested by numerical simulations (Eke et al. 1998).

3. RESULTS AND CONCLUSIONS

In Figure 1a we show the resulting best fit temperature functions using the power law approximation of Donahue and Voit (1999) and Blanchard et al. (1999) as we reconstructed them following their method. We used Horner et al. (1999)'s compilation of data taken from Edge et al. (1990) Henry and Arnaud (1991), and Markevitch (1998) (squares, diamonds, and triangles). The solid and dashed dotted lines represent best fit flat model, $\Omega_m = 0.27$, $\sigma_8 = 0.73$, $\alpha = 0.13$, of Donahue and Voit (they integrated over cluster formation epoch) and their best fit model, but not integrated over cluster formation epoch. Blanchard et al.'s best fit flat model, $\Omega_m = 0.735$, $\sigma_c = 0.623$, $\alpha = 0.18$, is represented by a long dashed line. The short dashed line represents Blanchard et al.'s best fit temperature function, but integrated over cluster formation epoch. Open model temperature functions (not shown) are very similar to those of the flat models. The difference between temperature functions from methods with and without integration over cluster formation epoch is significant even with existing data. The large difference between density parameters derived by Donahue and Voit, and Blanchard et al. is due to the different normalizations of the $M(T)$ relation and not integration over cluster formation, which can cause a change less than 0.1 in the density parameter (cf. next paragraph).

Figure 1b shows temperature functions of the best fit flat CDM model (best fit using method A) with $\Omega_m = 0.39$, $\Lambda = 0.61$, $h = 0.5$, and this best fit model, but temperature functions with and without taking the collapsed fraction and/or integrating over cluster formation epochs (our methods B, C and D). The data are the same as in figure 1a. The solid, long dashed, short dashed and dashed dot lines represent temperature functions using our methods A, B, C and D. Open models behave similarly so we do not show them here. The smallest effect is the correction for collapsed fraction of objects which is about the size of the error bars of the data, thus an important effect (compare methods A and B). A larger difference is caused by integration over formation epoch (methods A and C). If one fits models without integrating and/or taking collapsed fraction into account (methods B, C and D), one get matter densities $\Omega_m = 0.36$, $\Omega_m = 0.34$, and $\Omega_m = 0.31$ (assuming $h = 0.5$). Not taking either effects into account causes about 0.1 change in the derived matter density (compare results using methods A and D). As we can see, the temperature function is very sensitive to CDM model parameters if one uses a full CDM treatment, and thus a precise determination of the matter density is possible if the Hubble constant is known. We should keep in mind, however, that the normalization of the $M(T)$ relation, for example, causes much larger error in determining Ω_m . Since this method does not allow us to separate the density parameter from the Hubble constant, the result is a best fit function of the two. We can make use of the results from the power law approximation which gives the best fit density parameter, and check if the corresponding Hubble constant is reasonable using our methods. We find that our best fit CDM models yield the same matter




FIGURE 1. a) left panel: temperature functions using best fit power law approximations; b) right panel: temperature functions of the best fit CDM flat model using different methods. See text for details.

density as the best fit models of Donahue and Voit (1999) and Blanchard et al. (1999) if we use $h = 0.55$ and $h = 0.35$. Thus Blanchard et al.'s method is only marginally compatible to ours.

Figure 1b shows that systematic errors from interpretation of the temperature function are larger than the error bars on even the existing data. We conclude that, if the Hubble constant is known, a comparison between the observationally derived cluster temperature function and those derived from a full CDM treatment may yield an accurate determination of the density parameter (with an error less than 0.05), however, this can be done only if we find other ways to derive a correct theoretical model to interpret the data.

ACKNOWLEDGEMENTS

We thank N. Aghanim and R. Mushotzky for valuable discussions, and Don Horner for providing us his compilation of the temperature data. We thank N. Aghanim for her hospitality at the Institute of Astronomy of Orsay University.

REFERENCES

- Blanchard, A., et al.1999, astro-ph/9908037
- Donahue, M., and Voit, G. M., 1999, preprint, astro-ph/9907333
- Edge et al.1990, MNRAS, 245, 559
- Eke, V. R., Cole S., Frenk, C. S., and Henry, J. P., 1998, MNRAS, 298, 1145
- Horner, D. et al.1999, in preparation
- Hu, W, and Sugiyama, N, 1996, ApJ, 471, 542
- Markevitch, M., 1998, ApJ, 504, 27
- Martel, H., and Shapiro, P. R., 1999, astro-ph/9903425
- Henry, J. P., and Arnaud, K. A., 1991, ApJ, 372, 410
- Voges et al., 1999, A&A, 349, 389